ELSEVIER

Review of Palaeobotany and Palynology 92 (1996) 395

REVIEW OF PALAEOBOTANY AND PALYNOLOGY

Contents Volume 92, 1996

Research Papers: Taxonomy and palaeoecology of Early Cretaceous (Late Albian) angiosperm leaves from Alexander Island, Antarctica
D.J. Cantrill, G.J. Nichols 1
Fruits of Mastixioideae (Cornaceae) from the Paleogene of western North America
B.H. Tiffney, K.K. Haggard 29
Eocene *Nypa* from Regatta Point, Tasmania
M.S. Pole, M.K. Macphail 55
Spatial distribution of suspended pollen in the Mississippi River as an example of pollen transport in alluvial channels
A. Smirnov, G.L. Chmura, M.F. Lapointe 69
Charcoal in northcentral Alaskan lake sediments: relationships to fire and late-Quaternary vegetation history
C.J. Earle, L.B. Brubaker, P.M. Anderson 83
Neogene ecology of the Salto de Tequendama site (2475 m altitude, Cordillera Oriental, Columbia): the paleobotanical record of montane and lowland forests
V.M. Wijninga 97
A Pliocene *Podocarpus* forest mire from the area of the high plain of Bogotá (Cordillera Oriental, Colombia)
V.M. Wijninga 157
Anatomically preserved Permian *Noeggerathiopsis* leaves from east Antarctica
S. McLoughin, A.N. Drinnan 207
Nilssoniocladus in the Cretaceous Arctic: new species and biological insights
R.A. Spicer, A.B. Herman 229
A fertile pinnule fragment with spores of *Dicksonia* from Early Oligocene sediments in Tasmania
G.J. Jordan, M.K. Macphail, R.S. Hill 245
Pollen morphology of *Acanthopsis* Harvey, *Acanthus* L. and *Blepharis* Jussieu (Acanthaceae: Acantheae)
C.A. Furness 253
Ultrastructure of lower Paleozoic dyads from southern Ohio
W.A. Taylor 269
Preliminary palynological data on floristic and climatic changes during the Middle Eocene-Early Oligocene of the eastern Ebro Basin, northeast Spain
C. Cavagnetto, P. Anadón 281
Neogene environments in Australia, 1: re-evaluation of microfloras associated with important Early Pliocene marsupial remains at Grange Burn, southwest Victoria
M.K. Macphail 307
Palynology and paleobotany of the Early Pliocene section Río Frío 17 (Cordillera Oriental, Colombia): biostratigraphical and chronostratigraphical implications
V.M. Wijninga 329
Taxonomic stabilisation of dinoflagellate cyst taxa, as exemplified by two morphologically complex Early Cretaceous species
I.C. Harding 351
The presence of *Dinogymnium* (Dinophyceae) in Lower Cretaceous sediments from the northwest Tethys (southeast France and western Switzerland) and Gulf of Mexico areas: statigraphic and systematic consequences
L. Londeix, D. Pourtay, J.P.G. Fenton 367
Tianjinella, a new dinoflagellate genus from the Eocene of southern Tianjin, China
Changqnan He, Xuekun Sun 383
Technical Note: A simple method for microtome sectioning of prehistoric charcoal specimens, embedded in 2-hydroxyethyl methacrylate (HEMA)
A. Igersheim, O. Cichocki 389